



GOT THE CALL
you were exposed
to COVID-19?

MAKE THE CALL
about XOCOVA.



XOCOVA is the first-and-only prescription pill
used in people aged 12 years and older to help prevent
you from getting sick after exposure to COVID-19

COVID-19=coronavirus disease 2019.

What is XOCOVA?

XOCOVA (ensitrelvir) is a prescription medicine used to help prevent COVID-19 in adults and children aged 12 years and older following contact with a person who has COVID-19 (post-exposure prophylaxis). It is not known if XOCOVA is safe and effective in children under age 12.

Select Safety Information

Do not take XOCOVA if you are allergic to any of its ingredients or if you take certain medicines that may interact with XOCOVA and may cause severe or life-threatening side effects or death. Tell your healthcare provider (HCP) about all the medicines, vitamins and herbal supplements you take or plan to take, so your HCP can tell you if it is safe to take with XOCOVA. Tell your HCP if you are pregnant or plan to be, as XOCOVA may harm your unborn baby.

Please see Important Safety Information throughout and the full [Prescribing Information](#), including [Patient Information](#).

Be ready with a COVID-19 exposure plan

COVID-19 exposure can happen at any time.

Whether you're attending a crowded event like a wedding or concert, going on a family trip, or using public transit, be ready for when COVID-19 exposure happens.

Be prepared by proactively speaking with your healthcare professional about XOCOVA. That way you can ask for a prescription of XOCOVA to have on hand just in case you are exposed to COVID-19.



Don't let a
COVID-19 EXPOSURE CALL
catch you off guard

XOCOVA is used to help prevent COVID-19 in people aged 12 years and older after exposure.

COVID-19 can affect some people more seriously than others

COVID-19 can affect anyone, including individuals with certain risk factors who may be at a higher risk for COVID-related complications, such as those who have the following chronic health conditions or traits*:



Older adults



Obesity



Diabetes



Certain
organ conditions†



Prior or current
smoker



Weakened
immune system

*This is not an all-inclusive list of every condition that may increase risk of COVID-related complications. It is important to tell your healthcare professional about all your medical conditions.

†Includes heart, lung, and/or kidney conditions.

**Take action now
and ask about
XOCOVA**

**Act quickly after exposure to COVID-19
and ask your healthcare professional
about XOCOVA**

Important Safety Information

Do not take XOCOVA if you:

- are allergic to ensitrelvir or any of the ingredients in XOCOVA.
- take any of the following medicines, as these may interact with XOCOVA and may cause severe or life-threatening side effects or death:
 - apalutamide
 - carbamazepine
 - colchicine
 - dihydroergotamine
 - enzalutamide
 - eplerenone
 - ergotamine
 - finerenone
 - ivabradine
 - lomitapide
 - lumacaftor/ivacaftor
 - lurasidone
 - methylergonovine
 - phenytoin
 - pimozone
 - quinidine
 - rifampin
 - St John's wort
 - simvastatin
 - triazolam
 - voclosporin

These are not the only medicines that may cause serious or life-threatening side effects if taken with XOCOVA. XOCOVA may increase or decrease the levels of many medicines in your body. Tell your healthcare provider (HCP) about all the medicines you take because additional laboratory tests or dose changes of your other medicines

Important Safety Information (cont'd)

Do not take XOCOVA if you: (cont'd)

may be necessary during XOCOVA use. Your HCP may tell you about specific symptoms to watch for that may indicate you need to stop or decrease the dose of some of your other medicines.

Before taking XOCOVA, tell your HCP about all your medical conditions, including if you:

- are pregnant or plan to become pregnant. XOCOVA may harm your unborn baby. For females who are able to become pregnant, your HCP will check with you to see if you are pregnant before starting XOCOVA. Use effective birth control (contraception) during XOCOVA use and for 2 weeks after the final dose. Tell your HCP right away if you become pregnant or think you may be pregnant during XOCOVA use.

**Please see additional
Important Safety Information
throughout and the full
Prescribing Information,
including Patient Information.**



For people aged 12 years and older

XOCOVA was proven to help prevent COVID-19 after exposure

In a clinical study,
XOCOVA reduced the risk
of symptomatic COVID-19
after exposure by

67%*

If you were exposed to someone with COVID-19, XOCOVA can help reduce your risk of symptomatic COVID-19 if you start taking it right away within 3 days (72 hours) of exposure.

*Risk of symptomatic COVID-19 was assessed by Day 10 versus placebo.

Start XOCOVA right away after COVID-19 exposure


Take XOCOVA for 5 days

Take 3 pills together on Day 1, followed by 1 pill daily on Days 2 through 5.

Take XOCOVA at about the same time each day with or without food.

Do not stop taking XOCOVA without talking to your healthcare professional.

XOCOVA comes in a blister pack containing 7 pills. **Do not** remove XOCOVA pills from the blister pack until you are ready to take your dose.



Not actual pill size.



**Help reduce your risk
of getting sick after exposure
to COVID-19**

\$0

**Eligible commercially insured
patients may pay as little as
\$0 co-pay***

*Terms and Conditions apply.

Important Safety Information (cont'd)

Before taking XOCOVA, tell your HCP about all your medical conditions, including if you: (cont'd)

- are breastfeeding or plan to breastfeed. It is not known if XOCOVA passes into your breast milk. **Do not** breastfeed during XOCOVA use and for 2 weeks after the final dose.

Tell your HCP about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Your HCP can tell you if it is safe to take XOCOVA with other medicines. Do not start a new medicine without telling your HCP.

What are the possible side effects of XOCOVA?

XOCOVA may cause serious side effects, including:

- Allergic reactions, including severe allergic reactions (anaphylaxis). Stop taking XOCOVA and get medical help right away if you get any of the following allergic reaction symptoms: skin rash; hives; itching;

Important Safety Information (cont'd)

What are the possible side effects of XOCOVA? (cont'd)

swelling of the mouth, lips, tongue, or face; trouble swallowing or breathing; throat tightness; dizziness or feeling lightheaded; stomach-area (abdomen) pain; or vomiting.

The most common side effects of XOCOVA include: headache, diarrhea, and cough.

These are not all possible side effects of XOCOVA. For more info, ask your HCP or pharmacist. **You are encouraged to report negative side effects of prescription drugs to the FDA.** Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional
Important Safety Information
throughout and the full
[Prescribing Information](#),
including [Patient Information](#).

 **XOCOVA**
(ensitrelvir) 125 mg
tablets



Right after COVID-19 exposure, take action with XOCOVA



XOCOVA is the **first-and-only pill** used in people aged 12 years and older to help prevent you from getting sick* after exposure to COVID-19.



Take XOCOVA as soon as possible, **within 3 days** (72 hours) of being exposed to someone with COVID-19.

*In a clinical study, XOCOVA reduced the risk of symptomatic COVID-19 by 67% by Day 10 versus placebo (post-exposure prophylaxis).



Eligible commercially insured patients may pay \$0 co-pay[†]

[†]Terms and Conditions apply.

Visit [XOCOVA.com/faqs](https://xocova.com/faqs) to learn more about XOCOVA savings options

Take action. Plan ahead.

When you get the call, make the call.

Select Safety Information

Do not take XOCOVA if you are allergic to any of its ingredients or if you take certain medicines that may interact with XOCOVA and may cause severe or life-threatening side effects or death. Tell your healthcare provider (HCP) about all the medicines, vitamins and herbal supplements you take or plan to take, so your HCP can tell you if it is safe to take with XOCOVA. Tell your HCP if you are pregnant or plan to be, as XOCOVA may harm your unborn baby.

Please see Important Safety Information throughout and the full [Prescribing Information](#), including [Patient Information](#).

